

Orbactiv Order Form (ortavancin)

FAX TO: 817-472-7213

PATIENT INFORMATION				
Primary Language:	DOB: Allergies: on:			kg
<icd 10="" code="" require<="" td=""><td>DIAGNOSIS & CLI</td><td>NICAL INFORMATIO</td><td>N</td><td></td></icd>	DIAGNOSIS & CLI	NICAL INFORMATIO	N	
ICD 10 Code (PROVIDE COMPLETE CODE)				
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS</u> : Include culture report.				
PRESCRIPTION				
ORBACTIV (oritavancin)				
Start and follow each infusion with a 10 mL D5W flush DO NOT USE Normal Saline for dilution or flushing of IV line as it is incompatible with Orbactiv				
Loading Dose IV: Infuse 1200 mg in D5W for a total volume of 1000 mL as a single dose over 3 hours				
Is the patient on any other disease modifying therapy? Yes No If yes, please note therapy and last dose:				
Post Treatment Observations: The patient is observed for 30 minutes following the administration.				
Adverse Events: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol. Comments:				
PRESCRIBER INFORMATION				
Prescriber Name				
	NPI #:			
Address:	City:		State: Zip:	
Contact Name:	Phone:	Fax:	Email:	